

River at Pointe à Larrée, on the east coast ; rivers and fresh waters of the Cape of Good Hope.

16. *GERRES LINEOLATUS*, Günth.
17. *CHROMIS NILOTICUS*, Hasselq.
18. *ARIUS FALCARIUS*, Richards.
19. *LUTODEIRA CHANOS*, Forsk.
20. *ELOPS MACHNATA*, Forsk.
21. *ELOPS CYPRINOIDES*, Gm.
22. *ANGUILLA JOHANNÆ*, Günth.

4. Further Descriptions of New Species of Shells collected at Mauritius by Geoffrey Nevill, Esq. By HENRY ADAMS, F.L.S.

(Plate IV.)

*STOMATIA VARIEGATA*, H. Ad. (Pl. IV. fig. 1.)

*S. testa subrimata, ovato-oblonga, convexa, transversim costata, costis subdistantibus, interstitiis longitudinaliter striatis, rubra, albo variegata; spira prominula, apice acuto, sutura distincta; anfr. 3½, convexis; apertura ovali, labio calloso, labro supra ultimum anfractuum paulum ascende.*

Long. 9, lat. 7 mill.

Hab. Mauritius.

*DISCUS VORTICELLA*, H. Ad. (Pl. IV. figs. 2, 2a.)

*D. testa aperte umbilicata, subdiscoidea, tenui, oblique confertissime costulato-striata, fulva; spira vix elevata, sutura distincta; anfr. 4½, convexis, ultimo antice non descendente, superne subdepresso; umölico perspectivo, ½ diametri fere æquante; apertura obliqua, rotundato-lunari; perist. simplici, recto, marginibus conniventibus, columellari vix dilatato, dextro subsinuato.*

Diam. maj. 2¼, min. 2, alt. ¾ mill.

Hab. Bamboo, Mauritius.

*NANINA (ROTULA) CERNICA*, H. Ad. (Pl. IV. fig. 3.)

*N. testa imperforata, trochiformi-depressa, tenuissima, membranacea, oblique longitudinaliter striata, lineis spiralibus minutis decussata, corneo-lutescente; spira depresso-conica, apice obtusulo, sutura leviter impressa; anfr. 4, planatis, ultimo descendente, acute carinato, basi convexo; apertura diagonali,*



